



Chapter 1



2. The functional groups are identified below.



3. Amino acids, monosaccharides, nucleotides, and lipids are the four types of biological small molecules. Amino acids, monosaccharides, and nucleotides can form polymers of proteins, polysaccharides, and nucleic acids, respectively.

4. (a) *N*-acetylglucosamine is a monosaccharide.

(b) CMP is a nucleotide.

(c) Homocysteine is an amino acid.

(d) Cholesteryl ester is a lipid.

5. (a) Mainly C and H plus some O

(b) C, H, and O

(c) C, H, O, and N plus small amounts of S

6. It is a lipid (it is actually lecithin). It is mostly C and H, with relatively little O and only one N and one P. It has too little O to be a carbohydrate, too little N to be a protein, and too little P to be a nucleic acid.

7. (a) You should measure the nitrogen content, since this would indicate the presence of protein (neither lipids nor carbohydrates contain nitrogen).

(b) You could add the compound that contains the most nitrogen, compound B, which is melamine. (Melamine is a substance that has been


added to some pet foods and milk products from China so that they would appear to contain more protein. Melamine is toxic to pets and children.)

(c) Compound C is an amino acid, so it would already be present in protein-containing food.

8. A diet high in protein results in a high urea concentration, since urea is the body's method of ridding itself of extra nitrogen. Nitrogen is found in proteins but is not found in significant amounts in lipids or carbohydrates. A low-protein diet provides the patient with just enough protein for tissue repair and growth. In the absence of excess protein consumption, urea production decreases, and this puts less strain on the patient's weakened kidneys.



9. All amino acids have carboxylate groups. All have primary amino groups except for proline, which has a secondary amino group.

10. The carbon marked with an asterisk is chiral. This means that alanine has two possible enantiomers, or mirror-image isomers.



11. Asn has an amido group and Cys has a sulfhydryl group.

12. Two hydrogen atoms and one oxygen atom are lost when Asn and Cys form a dipeptide. This is an example of a condensation reaction. The carboxylate functional group on the Asn is lost and the amino functional group on the Cys is lost. An amido group is formed.



14. (a) Fructose has the same molecular formula, $\text{C}_6\text{H}_{12}\text{O}_6$, as glucose.

(b) Fructose is a ketone, whereas glucose is an aldehyde.

15. Uracil has a carbonyl functional group, whereas cytosine has an amino functional group.

16. Nucleotides consist of a five-carbon sugar, a nitrogenous ring, and one or more phosphate groups linked covalently together.

17. As described in the text, palmitate and cholesterol are highly nonpolar and are therefore insoluble in water. Both are highly aliphatic. Alanine is water soluble because its amino group and carboxylate group are ionized, which render the molecule "saltlike." Glucose is also water soluble because its aldehyde group and many hydroxyl groups are able to form hydrogen bonds with water.

2 Ch 1 Solutions

18. Glucose has several hydroxyl groups and is a polar molecule. As such, it will have difficulty crossing the nonpolar membrane. The 2,4-dinitrophenol molecule consists of a substituted benzene ring and has greater nonpolar character. Of the two molecules, the 2,4-dinitrophenol will traverse the membrane more easily.

19. DNA forms a more regular structure because DNA consists of only four different nucleotides, whereas proteins are made up of as many as 20 different amino acids. In addition, the 20 amino acids have much more individual variation in their structures than do the four nucleotides. Both of these factors result in a more regular structure for DNA. The cellular role of DNA relies on the *sequence* of the nucleotides that make up the DNA, not on the overall shape of the DNA molecule itself. Proteins, on the other hand, fold into unique shapes, as illustrated by endothelin in Figure 1-4. The ability of proteins to fold into a wide variety of shapes means that proteins can also serve a wide variety of biochemical roles in the cell. According to Table 1-2, the major roles of proteins in the cell are to carry out metabolic reactions and to support cellular structures.

20. Polysaccharides serve as fuel-storage molecules and can also serve as structural support for the cell.

21. The pancreatic amylase is unable to digest the glycosidic bonds that link the glucose residues together in cellulose. Figure 1-6 shows the structural differences between starch and cellulose. Pancreatic amylase binds to starch prior to catalyzing the hydrolysis of the glycosidic bond; thus, the enzyme and the starch must have shapes that are complementary. The enzyme would be unable to bind to the cellulose, whose structure is much different from that of starch.

22. Cellulose cannot be digested in mammals and therefore the energy yield is 0 kilocalories per gram. Although both starch and glycogen are polymers of glucose, the glucose residues are linked together differently in the two molecules and pancreatic amylase is unable to hydrolyze the glycosidic bonds (see Solution 21). Cellulose provides no energy to the diet but is an important component of the diet as fiber.

23. A positive entropy change indicates that the system has become more disordered; a negative entropy change indicates that the system has become more ordered.

- | | |
|--------------|--------------|
| (a) negative | (b) positive |
| (c) positive | (d) positive |
| (e) negative | |

- 24.** (a) decrease
(b) increase

25. The polymeric molecule is more ordered and thus has less entropy. A mixture of constituent monomers has a large number of different arrangements (like the balls scattered on a pool table) and thus has greater entropy.

26. Entropy increases as the reactants (7 molecules) are converted to products (12 molecules).

27. The dissolution of ammonium nitrate in water is a highly endothermic process, as indicated by the positive value of ΔH . This means that when ammonium nitrate dissolves in water, the system absorbs heat from the surroundings and the surroundings become cold. The plastic bag containing the ammonium nitrate becomes cold and can be used as a cold pack to treat an injury.

28. The dissolution of calcium chloride in water is a highly exothermic process, as indicated by the negative value of ΔH . This means that when calcium chloride dissolves in water, the system loses heat to the surroundings and the surroundings become warm. The plastic bag holding the calcium chloride dissolution becomes warm and can be used as a hot pack by the hiker at cold temperatures.

29. First, calculate ΔH and ΔS , as described in Sample Calculation 1-1:

$$\begin{aligned}\Delta H &= H_B - H_A \\ \Delta H &= 60 \text{ kJ} \cdot \text{mol}^{-1} - 54 \text{ kJ} \cdot \text{mol}^{-1} \\ \Delta H &= 6 \text{ kJ} \cdot \text{mol}^{-1} \\ \Delta S &= S_B - S_A \\ \Delta H &= 43 \text{ J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1} - 22 \text{ J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1} \\ \Delta H &= 21 \text{ J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}\end{aligned}$$

- (a) $\Delta G = (6000 \text{ J} \cdot \text{mol}^{-1}) - (4 + 273 \text{ K})(21 \text{ J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1})$
 $\Delta G = 180 \text{ J} \cdot \text{mol}^{-1}$

The reaction is not favorable at 4°C.

(b) $\Delta G = (6000 \text{ J} \cdot \text{mol}^{-1}) - (37 + 273 \text{ K})(21 \text{ J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1})$
 $\Delta G = -510 \text{ J} \cdot \text{mol}^{-1}$

The reaction is favorable at 37°C.

- 30.** $0 > 15,000 \text{ J} \cdot \text{mol}^{-1} - (T)(51 \text{ J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1})$
 $-15,000 > -(T)(51 \text{ K}^{-1})$
 $15,000 < (T)(51 \text{ K}^{-1})$
 $294 \text{ K} < T$

The reaction is favorable at temperatures of 21°C and higher.

- 31.** $0 > -14.3 \text{ kJ} \cdot \text{mol}^{-1} - (273 + 25 \text{ K})(\Delta S)$
 $14.3 \text{ kJ} \cdot \text{mol}^{-1} > -(273 + 25 \text{ K})(\Delta S)$
 $-48 \text{ J} \cdot \text{mol}^{-1} > \Delta S$
 ΔS could be any positive value, or it could have a negative value smaller than $-48 \text{ J} \cdot \text{mol}^{-1}$.
- 32.** $-63 \text{ kJ} \cdot \text{mol}^{-1} = \Delta H - (273 + 25 \text{ K})(190 \text{ J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1})$
 $\Delta H = -63 \text{ kJ} \cdot \text{mol}^{-1} + 56.6 \text{ kJ} \cdot \text{mol}^{-1}$
 $\Delta H = -6.4 \text{ kJ} \cdot \text{mol}^{-1}$

The reaction is exothermic.


- 33.** Process (d) is never spontaneous.
- 34.** (a) Because the rubber band becomes cooler, it is absorbing heat. Therefore, its change in enthalpy is positive.
(b) In order for ΔG to be negative (a spontaneous process), the entropy change must be negative so that it can compensate for an unfavorable (positive) change in enthalpy.
- 35.** The dissolution of urea in water is an endothermic process and has a positive ΔH value. In order to be spontaneous, the process must also have a positive ΔS value in order for the free energy change of the process to be negative. Solutions have a higher order of entropy than the solvent and solute alone.

- 36.** (a) The reaction is exothermic because the value of ΔH is negative.
(b) $\Delta G = \Delta H - T\Delta S$
 $-17200 \text{ J} \cdot \text{mol}^{-1} = -9500 \text{ J} \cdot \text{mol}^{-1} - (310 \text{ K})(\Delta S)$
 $\Delta S = 25 \text{ J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$

The positive value of ΔS indicates that the reaction proceeds with an increase in entropy.


(c) The ΔH term makes a greater contribution to the ΔG value. This indicates that the reaction is spontaneous largely because the reaction is exothermic.

- 37.** (a) The conversion of glucose to glucose-6-phosphate is not favorable because the ΔG value for the reaction is positive, indicating an endergonic process.
(b) If the two reactions are coupled, the overall reaction would be the sum of the two individual reactions. The ΔG value would be the sum of the ΔG values for the two individual reactions.



Coupling the conversion of glucose to glucose-6-phosphate with the hydrolysis of ATP has converted an unfavorable reaction to a favorable reaction. The ΔG value of the coupled reaction is negative, which indicates that the reaction as written is favorable.


- 38.** (a) The reaction is not favorable because the ΔG value for the reaction is positive, indicating an endergonic process.
(b)



The coupled reaction is spontaneous because the ΔG value is negative.

- 39.** C (most oxidized), A, B (most reduced)
- 40.** (a) reduction
(b) oxidation


41. (a) oxidized (b) oxidized
(c) oxidized (d) reduced
42. (a) oxidizing agent (b) oxidizing agent
(c) oxidizing agent (d) reducing agent
43. (a) Palmitate's carbon atoms, which have the formula $\text{—CH}_2\text{—}$, are more reduced than CO_2 , so their reoxidation to CO_2 can release free energy.
(b) Because the $\text{—CH}_2\text{—}$ groups of palmitate are more reduced than those of glucose (—HCOH—), their conversion to the fully oxidized CO_2 would be even more thermodynamically favorable (have a larger negative value of ΔG) than the conversion of glucose carbons to CO_2 . Therefore, palmitate carbons could provide more free energy than glucose carbons.
44. The complete oxidation of stearate to CO_2 yields more energy because 17 of the 18 carbons of stearate are fully reduced. The conversion of these carbons to CO_2 provides more free energy than some of the carbons of α -linolenate, which participate in double bonds and are therefore already partially oxidized.
45. The experiment was significant because it demonstrated that it was possible to synthesize the building blocks of biological macromolecules (amino acids, carbohydrates, and nucleic acids) using only inorganic gases as starting materials and lightning as an energy source, that is, the conditions that most likely existed in the prebiotic world.
46. The first biological molecules would have had to polymerize, and they would have had to find some way to make copies of themselves.
47. Morphological differences, which are useful for classifying large organisms, are not useful for bacteria, which often look alike. Furthermore, microscopic organisms do not leave an easily interpreted imprint in the fossil record, as vertebrates do. Thus, molecular information is often the only means for tracing the evolutionary history of bacteria.
48. It is difficult to envision how a single engulfment event could have given rise to a stable and heritable association of the eukaryotic host and the bacterial dependent within a single generation. It is much more likely that natural selection gradually promoted the interdependence of the cells. Over many generations, genetic information supporting the association would have become widespread.
49. (a) H15 and H7 are closely related, as are H4 and H14.
(b) H4 and H14 are most closely related to H3




Chapter 2

1. The water molecule is not perfectly tetrahedral because the electrons in the nonbonding orbitals repel the electrons in the bonding orbitals more than the bonding electrons repel each other. The angle between the bonding orbitals is therefore slightly less than 109° .

2. Because the partial negative charges are arranged symmetrically (and the shape of the molecule is linear), the molecule as a whole is not polar.




3. Ammonia is polar because it has one unshared electron pair. Its shape is trigonal pyramidal, and the molecule is not symmetrical. Nitrogen is more electronegative than hydrogen, so partial negative charges reside on the nitrogen and partial positive charges on the hydrogens.




4. Water has the highest melting point because each water molecule forms hydrogen bonds with four neighboring water molecules, and hydrogen bonds are among the strongest intermolecular forces. Ammonia is also capable of forming hydrogen bonds, but they are not as strong (due to the electronegativity difference between hydrogen and nitrogen). Methane cannot form hydrogen bonds; the molecules are attracted to their neighbors only via weak London dispersion forces.

5. The arrows point toward hydrogen acceptors and away from hydrogen donors.



6. Compound A does not form hydrogen bonds (the molecule has a hydrogen bond acceptor but no hydrogen bond donor). Compounds B and C form hydrogen bonds as shown because each molecule contains at least one hydrogen bond donor and a hydrogen bond acceptor. The molecules in D do not form hydrogen bonds with each other, whereas the molecules in E do because ammonia has a hydrogen bond donor and diethyl ether has a hydrogen bond acceptor.



7. (a) van der Waals forces (dipole-dipole interactions)
(b) Hydrogen bonding
(c) van der Waals forces (London dispersion forces)
(d) Ionic interactions

8. (a) $\text{H} < \text{C} < \text{N} < \text{O} < \text{F}$
(b) The greater an atom's electronegativity, the more polar its bond with H and the greater its ability to act as a hydrogen bond acceptor. Thus, N, O, and F, which have relatively high electronegativity, can form hydrogen bonds, whereas C, whose electronegativity is only slightly greater than hydrogen's, cannot.

9. From the highest melting point to the lowest melting point: C, B, E, A, D. Compound C (urea, melting point 133°C) has three functional groups that can serve as hydrogen donors and/or acceptors. Compound B (acetamide, melting